

Deep $g'r'i'z'$ GMOS Imaging of the Dwarf Irregular Galaxy Kar 50¹

T. J. Davidge

Canadian Gemini Office, Herzberg Institute of Astrophysics,
National Research Council of Canada, 5071 West Saanich Road,
Victoria, B.C. Canada V9E 2E7
email: tim.davidge@nrc.ca

ABSTRACT

Images obtained with the Gemini Multi-Object Spectrograph (GMOS) are used to investigate the stellar content and distance of the dwarf irregular galaxy Kar 50. The brightest object is an HII region, and the bright stellar content is dominated by stars with $g' - r' < 0$. The tips of the main sequence and the red giant branch are tentatively identified near $r' = 24.9$ and $i' \sim 25.5$, respectively. The galaxy has a blue integrated color with no significant color gradient, and we conclude that Kar 50 has experienced a recent galaxy-wide episode of star formation. The distance estimated from the brightest blue stars indicates that Kar 50 is behind the M81 group, and this is consistent with the tentative RGB-tip brightness. Kar 50 has a remarkably flat central surface brightness profile, even at wavelengths approaching $1\mu\text{m}$, although there is no evidence of a bar. In the absence of another large star-forming episode, Kar 50 will evolve into a very low surface brightness galaxy.

Subject headings: galaxies: individual (Kar 50) - galaxies: dwarf - galaxies: stellar content

1. INTRODUCTION

The Local Group contains over 30 dwarf galaxies that span a range of morphological types (e.g. van den Bergh 2000). While such nearby systems are fundamental laboratories

¹Based on observations obtained at the Gemini Observatory, which is operated by the Association of Universities for Research in Astronomy, Inc., under a co-operative agreement with the NSF on behalf of the Gemini partnership: the National Science Foundation (United States), the Particle Physics and Astronomy Research Council (United Kingdom), the National Research Council of Canada (Canada), CONICYT (Chile), the Australian Research Council (Australia), CNPq (Brazil), and CONICET (Argentina).

for investigating dwarf galaxy evolution, many are satellites of M31 and the Milky-Way, and hence have been affected by hierarchical interactions (e.g. Mayer et al. 2001). In order to develop a more comprehensive understanding of dwarf galaxies, including the effects of environment, it is necessary to study objects outside of the Local Group.

Located at a distance of ~ 3.5 Mpc, the M81 group contains a number of dwarf galaxies, some of which are systematically different from those in the Local Group (Caldwell et al. 1998). Karachentseva, Karachentsev & Börngen (1985, hereafter KKR) list possible dwarf members of the M81 group. Many of these have yet to be studied in detail, and some will likely turn out not to be members of the M81 group. This being said, the census of galaxies in the M81 group is likely far from complete, and Froebrich & Meusinger (2000) report the detection of additional possible low surface brightness members.

One of the faintest, and hence intrinsically most interesting, systems in the KKR compilation is Kar 50. KKR noted that Kar 50 is ‘unusually filamentary with barely outlined knots’, and they partially resolved Kar 50 into stars using images recorded during 1 arcsec seeing conditions. Various observational properties of Kar 50 are summarized in Table 1.

Motivated by the possible intrinsic faintness of Kar 50, coupled with its unusual appearance, we decided to include this galaxy in a background field obtained as part of a program to study the outer regions of NGC 2403. The observations and the data reduction procedures are discussed in §2, and these data are used to investigate the integrated photometric properties of Kar 50 in §3. Our i' and z' images are of particular interest for work of this nature, as they sample redder wavelengths than previous studies, and hence are better probes of any cool stellar component, such as might arise from an old population. The properties of the brightest stars in Kar 50 are investigated in §4. A distance based on the brightest stars is calculated in §5, and we conclude that Kar 50 is located well behind the M81 group. A brief summary and discussion of the results follows in §6.

2. OBSERVATIONS & REDUCTIONS

The data were recorded with the Gemini Multi-Object Spectrograph (GMOS) on the Gemini North telescope during 2 nights in November 2001 as part of a system verification program to investigate the outer regions of NGC 2403 (Davidge & Roy 2002, in preparation). Readers are referred to Crampton et al. (2000) for a detailed description of GMOS. The GMOS detector consists of three butttable 2048×4608 EEV CCDs, with each pixel subtending 0.0727 arcsec on a side. The sky field has a 5.5 arcmin diameter, although

the detector covers a much larger area so that spectra of objects near the edge of the sky field can be obtained.

Details of the observations, including dates, exposure times, and image quality are listed in Table 2. The sky conditions were photometric when the data were obtained, and standard stars were observed on each night. The CCDs were binned 2×2 during readout to better match the seeing conditions, so that each superpixel covers 0.145 arcsec on a side. Bias frames and twilight flats were also obtained on each night.

The data were reduced using tasks in the Gemini IRAF package. The reduction pipeline included (1) removal of CCD-to-CCD gain differences, (2) subtraction of the bias pattern, (3) division by a twilight sky flat, and (4) mosaicking of the processed CCD output to form a single image. Interference fringes were removed from the i' and z' data by subtracting a fringe frame that was created by combining observations of different fields. The processed images were then aligned and median-combined on a filter-by-filter basis to reject cosmic rays.

3. THE APPEARANCE AND INTEGRATED PHOTOMETRIC PROPERTIES OF KAR 50

The portion of the final r' image containing Kar 50 is shown in Figure 1. These data have a finer angular resolution and go deeper than previous observations of Kar 50. There is an absence of structures such as a bar or a central nucleus. Concentrations of stars are evident in the outer envelope, and these are likely the ‘filamentary’ features referred to by KKR. There are a number of stars evident throughout the galaxy, although the density of sources outside of the main body of the galaxy is significant, and it can be anticipated that many of the objects detected in Kar 50 may be foreground stars and background galaxies.

Karachentseva et al. (1987; hereafter K87) included Kar 50 in their investigation of the structural properties of M81 group dwarf galaxies. Kar 50 is the most compact object in their sample, and falls within the scatter envelope of the relations between size, surface brightness, and total intrinsic brightness defined by low surface brightness dwarf spheroidals. Both K87 and Barazza, Binggeli, & Prugniel (2001, hereafter BBP) find that the central surface brightness profile of Kar 50 at visible and red wavelengths is very flat, changing by only a few tenths of a magnitude per square arcsec within a few arcsec of the galaxy center. The surface brightness profile of the halo follows an exponential relation (BBP).

A series of aperture measurements, positioned on the geometric center of Kar 50 as

established from the i' image, were made in each filter. A technique similar to that employed by Davidge (1992) and Piotto, King, & Djorgovski (1988) to investigate the color profiles of globular clusters was used to suppress bright resolved stars. In particular, successive annuli were divided into six azimuthal segments, and the median surface brightness within each group of segments was assigned to that annulus.

The resulting g' , r' , i' and z' surface brightness profiles are shown in Figure 2. The error bars show the uncertainty in the surface brightness measurements in the last annulus due to uncertainties in the local sky level, which were estimated by examining the background at various locations in the vicinity of Kar 50. The light profiles are remarkably flat near the galaxy center, with a scatter of only ± 0.04 mag in each filter in the central 8 arcsec. Light profiles obtained without azimuthal median filtering are similar to those in Figure 2.

The light profiles measured in i' and z' are of particular interest, as previous studies of Kar 50 have not sampled wavelengths longward of $\sim 0.7\mu\text{m}$. Not only are the i' and z' data less susceptible to reddening than the g' and r' observations, but they also are a more sensitive probe of any underlying old and/or intermediate-age population. It is evident from Figure 2 that there are no hints of a central concentration in the main body of the galaxy, even in i' and z' .

The $g' - i'$ and $r - z'$ color profiles of Kar 50 are shown in Figure 3. The color profiles do not show significant gradients when $r < 15$ arcsec, and the only deviations from a flat color profile occur at larger radii, where the uncertainties in the color measurements become significant. The onset of the exponential light profile near 8 arcsec is not accompanied by an obvious change in stellar content. The stellar content of Kar 50 is thus well mixed throughout the galaxy.

Lenz et al. (1998) investigate the photometric properties of stars in the SDSS filter system, and colors predicted from their $\log(g) = 4.5$ solar metallicity sequences for three effective temperatures are marked in Figure 3. There is a tendency for the observations at longer wavelengths to sample cooler populations, as expected for a composite stellar system. Nevertheless, the relatively blue ($r' - z'$) colors indicate that the integrated light from Kar 50 is dominated by hot young stars, even near $1\mu\text{m}$.

4. THE RESOLVED STELLAR CONTENT

The current investigation is restricted to objects within ~ 30 arcsec of the center of Kar 50, and photometry of sources over the remainder of the GMOS field of view will be discussed in a future paper dealing with the outer regions of NGC 2403 (Davidge

& Roy 2002, in preparation). The brightnesses of individual stars were measured with the PSF-fitting routine ALLSTAR (Stetson & Harris 1988), using co-ordinates, initial brightnesses, and PSFs obtained from DAOPHOT (Stetson 1987) tasks. The PSFs were constructed from moderately bright stars located over the entire GMOS field of view.

The $(r', g' - r')$ and $(i', r' - i')$ CMDs of stars in two radial intervals centered on Kar 50 are shown in Figures 4 and 5; we do not consider the z' measurements here as they do not go as deep as the other data. The CMDs labelled ‘Kar 50’ contain sources within 22 arcsec of the galaxy center, while the CMDs labelled ‘background’ consist of sources between 22 to 31 arcsec from the galaxy center; the former region has the same area as the latter. The radial boundaries used to construct the CMDs are marked in Figure 1.

The number density of the brightest stars in the left hand panels of Figure 4 and 5 is only slightly higher than that of background objects having comparable brightness. At fainter levels the contrast with respect to background sources increases, and there is a concentration of stars in the Kar 50 CMDs with relatively blue colors ($g' - r' \sim -0.4$, and $r' - i' \sim 0$) near $r' \sim i' \sim 25$. Based on the blue colors, we identify this clump of stars as the onset of the upper main sequence. The r' and i' observations probe a redder population than the g' data, and there is a clump of stars near the faint end of the $(i', r' - i')$ CMD of Kar 50 at $i' \sim 25.5$ and $r' - i' \sim 0.3$. This color is consistent with these objects being red giants, and we tentatively identify this collection of stars as the RGB-tip. The identifications of the main sequence and RGB are discussed further in §6.

Contamination from foreground stars and background galaxies complicates efforts to study stars brighter than $r' \sim i' = 25$ in Figures 4 and 5, but color information can be used to identify a sample of bright stars that likely belong to Kar 50. Stars occupy a distinct region of the $(g' - r', r' - i')$ two-color diagram (TCD) (e.g. Krisciunas, Margon, & Szkody 1998), and multi-color observations can be used to identify non-stellar interlopers (e.g. Fan 1999). The $(g' - r', r' - i')$ TCDs of the Kar 50 and background fields are compared in Figure 6; the majority of objects plotted in this figure have $r' < 25.5$. The solid line shows the predicted locus of solar-metallicity $\log(g) = 4.5$ stars from Table 4 of Lenz et al. (1998), while the dashed line is the ridgeline of the SDSS stellar sequence from Figure 3 of Schneider et al. (2001); the data points in Figure 6 agree better with the SDSS sequence than the model sequence.

The objects within 22 arcsec of Kar 50 plotted in Figure 6 tend to be bluer than those in the background, in the sense that the majority of sources in Kar 50 have $(g' - r') < 0$, whereas only two objects in the background region are this blue. Given this obvious difference, it was assumed that all sources with $g' - r' < 0$ within 22 arcsec of Kar 50 belong to the galaxy, while the sources with $g' - r' > 0$ are likely contaminating objects. This

criterion for Kar 50 membership is further verified by comparing the number of objects with $g' - r' > 0$ in the right and left hand panels of Figure 6: there are 14 sources with $(g' - r') > 0$ within 22 arcsec of the galaxy center, and 15 such objects in the background field. This excellent agreement indicates that Kar 50 likely does not harbour a substantial component with $r' < 25$ and $(g' - r') > 0$; rather, hot stars appear to dominate the bright stellar content of Kar 50.

The $(r', g' - r')$ and $(i', r' - i')$ CMDs of Kar 50, corrected for line of sight absorption using $E(B - V)$ determined from the Schlegel, Finkbeiner, & Davis (1998) maps, are shown in Figure 7. The imposition of a selection criterion based on $g' - r'$ has a major impact on the interpretation of the CMDs when $r' < 25$, in that many of the sources that might have been selected as candidate bright members of Kar 50 in the absence of $g' - r'$ color information likely do not belong to the galaxy; the impact is most dramatic on the $(i', r' - i')$ CMD, where sources with $g' - r' > 0$ are indicated. The hot stars in Kar 50 with $r_0 \sim 24.8$ that we associate with the upper main sequence have $(g' - r')_0 \leq -0.4$. The solar metallicity models of Lenz et al. (1998) predict that the blue limit for stars is $g' - r' \sim -0.5$, and so it appears that upper main sequence stars in Kar 50 are not subject to internal reddening in excess of $E(g' - r') = 0.1$ mag.

The brightest blue object in the $(r', g' - r')$ CMD of Kar 50, the location of which is indicated in Figure 1, has $r' = 22.7$ and $g' - r' = -0.3$. While the $g' - r'$ color is within the acceptable range for stellar sources, this object has an extremely blue $r' - i$ color, with $r' - i' \sim -0.9$, placing it in a region of the $(g' - r', r' - i')$ TCD that is populated by compact emission line galaxies (Fan 1999). This object is likely an HII region in Kar 50, with the g' and r' measurements dominated by strong [OIII] and H α emission, respectively. This object is not extended in the 0.6 arcsec FWHM r' and i' images, indicating that it has a linear size < 40 parsecs using the distance derived in the next Section. As the only large HII region in Kar 50, this object has the potential of providing unique information about the gas-phase chemical composition of the galaxy.

5. THE DISTANCE TO KAR 50

The integrated colors and stellar content of Kar 50 indicate that the galaxy has experienced recent star formation, and so the properties of the brightest blue stars provide a means of estimating the distance to this system. Rozanski & Rowan-Robinson (1994) discuss the brightest stars as distance indicators, and give a relation between the mean magnitude of the three brightest blue stars and galaxy brightness. The calibration in their Figure 10c has ± 1 mag scatter, largely because the bright blue stellar content (1) is subject

to stochastic effects, (2) is sensitive to the recent star-forming history, and (3) may be affected by internal extinction. Despite the large scatter in the calibration, the brightest blue stars still provide interesting limits on the distance of Kar 50, which are consistent with other indicators (§6).

The brightnesses and colors of the three brightest objects with stellar spectral-energy distributions (SEDs) in Kar 50 are summarized in Table 3. The V_0 and $(B - V)_0$ entries for each star are in the AB₉₅ system, and these were computed from the transformation equations given in Section 2.1 of Krisciunas et al. (1998). We prefer the Krisciunas et al (1998) transformation relations over those given by Fukugita et al. (1996) as the former are based on actual stellar observations; nevertheless, a potential source of uncertainty is that the brightest stars in Kar 50 are supergiants, whereas the transformation relations are likely dominated by main sequence and giant branch stars. Adopting $B_{AB95} = -0.12$ for Vega (Fukugita et al. 1996) then $B_0(3) = 23.3$ in the Johnson system.

The relation between $M_B(3)$ and M_{B_T} in Figure 10c of Rozanski & Rowan-Robinson (1994) was iterated until convergence, and the Kar 50 distance modulus was found to be 30.4; the integrated brightness of Kar 50, computed from the B_T value in Table 1, is then $M_B = -12.7$. The uncertainty in these quantities is dominated by the scatter in the calibrating relation, and so we assign ± 1 mag errors to the distance modulus and M_B .

6. DISCUSSION & SUMMARY

Deep multicolor images obtained with the GMOS on the Gemini North telescope have been used to investigate the structure and stellar content of the dwarf irregular galaxy Kar 50. The presence of a significant population of bright blue stars, coupled with the blue integrated colors of the galaxy and the flat color profiles, which indicate that the stellar content of the galaxy is well mixed, indicate that Kar 50 has experienced a recent galaxy-wide star-forming episode. Kar 50 may thus be one of the small percentage of dwarf irregular galaxies that at any given time host a strong burst of star formation (Dohm-Palmer et al. 1998). In the remainder of this Section we discuss (1) the stellar content of Kar 50, with emphasis on checking the distance estimated from the brightest blue stars, and (2) the evolution of the galaxy.

6.1. The Distance and Stellar Content of Kar 50

While contamination from foreground stars and background galaxies complicates efforts to study the resolved stellar content of Kar 50 when $r' < 25$, in §4 it is demonstrated that a secure population of bright galaxy members can be identified using $g' - r'$ color. The application of a color criterion to reject contaminating objects will of course cause some objects, in this case those that have red colors, to be excluded from the analysis. The brightest red supergiants (RSG)s in dwarf galaxies have $M_V \sim -7$ (Rozanski & Rowan-Robinson 1994), and thus would be expected to occur near $r' \sim 23.7$ at the distance of Kar 50. While this is well within the detection limits of our data, the agreement between the numbers of objects with $g' - r' < 0$ in the Kar 50 and background fields indicates that RSGs do not occur in great numbers in Kar 50.

The brightest object in Kar 50 has an SED suggesting that it is an HII region, and if this is the case then the r' flux will be dominated by $H\alpha$ emission. With an assumed distance modulus of 30.4, this object has a luminosity of 1.3×10^{37} ergs/sec in r' , which is consistent with the peak $H\alpha$ luminosity of HII regions in other irregular galaxies (Youngblood & Hunter 1999). With a distance modulus of 27.5 then the luminosity of the HII region drops to 8.9×10^{35} erg/sec, which is near the lower end of what is seen in irregular galaxies. Thus, the luminosity of this object is consistent with it being an HII region. Nevertheless, this object is very compact, as the maximum size of 40 parsecs is a factor of 2 smaller than what is seen among HII regions in other irregular galaxies (Youngblood & Hunter 1999).

The distance of 12^{+7}_{-4} Mpc measured from the brightest blue stars places Kar 50 well behind the M81 group. There are significant uncertainties inherent to the use of the brightest blue stars as standard candles, and so in the remainder of this section we check if this distance is consistent with the properties of other resolved objects in Kar 50.

The RGB-tip is a prime distance indicator for old stellar systems, as it produces a clear break in luminosity functions and CMDs. The RGB-tip occurs near $M_{i'} \sim -4$ in old metal-poor populations (e.g. Davidge et al. 2002), and hence would occur at $i' \sim 23.5$ if Kar 50 were in the M81 group. Given the evidence for moderately recent star formation, a well populated AGB sequence might also be anticipated above the RGB-tip in Kar 50. There is no evidence in the CMDs of Kar 50 of either a well-defined RGB peaking near $i' \sim 23.5$ or a statistically significant AGB sequence when $i' < 23.5$, which would produce an excess red population with respect to the background at this brightness. Moreover, it is evident from Figure 7 that many of the sources on the $(i', r' - i')$ CMD have $(r' - i') \sim 0.0$, whereas RGB stars will have $r' - i' \geq 0.3$. There is an apparent onset of stars in the $(i', r' - i')$ CMD with $i' \sim 25.4$ and $r' - i' \sim 0.3$. If this is the RGB-tip then this supports a distance modulus ~ 29.4 , which falls within the ± 1 mag errors in the distance modulus computed

from the brightest blue stars. We are reluctant to assign a firm distance estimate based on the RGB-tip because it is at the faint limit of our data, and AGB stars can complicate distances measured from this feature.

The brightest stars at visual wavelengths in star-forming galaxies tend to have spectral types A – F (e.g. Humphreys & McElroy 1984), and the stars used to estimate the distance to Kar 50 have colors that are consistent with this range of spectral types (§5). Humphreys & McElroy (1984) list the effective temperatures and brightnesses of supergiants and upper main sequence stars in the Milky-Way and Magellanic Clouds, and we have used these data to locate these sequences on the $(r', g' - r')$ CMD. Colors were calculated using the relation between effective temperature and $g' - r'$ predicted by the solar metallicity $\log(g)=4.5$ atmosphere models of Lenz et al. (1998), while $M_{r'}$ was computed from the Krisciunas et al. (1998) transformation relations. The Lenz et al. (1998) models have surface gravities appropriate for main sequence stars, but not supergiants, and this likely introduces systematic errors in $g' - r'$ on the order of a few hundredths of a magnitude in the placement of the supergiant sequence.

The loci defined by Ia and Ib supergiants, as well as the main sequence, are shown in Figure 8 based on the adopted distance modulus of 30.4. The type Ia supergiant sequence loosely matches the upper envelope of stars in Kar 50, while the Ib sequence passes through the main body of the CMD. In addition, the bright end of the main sequence is in reasonable agreement with the clump of stars we identified in §4 as the upper main sequence tip. Based on these comparisons we predict that spectroscopic studies will reveal that 1) the brightest stars in Kar 50 are Ia supergiants, and (2) the brightest main sequence stars in Kar 50 are early-type O stars.

6.2. Some Comments on the Structure and Evolution of Kar 50

The light and color profiles of Kar 50 offer clues about the past evolution of the galaxy. The light profile of Kar 50 is flat over a linear scale of ~ 1 kpc, while the color profile suggests that the stellar content is well mixed throughout the galaxy. Light and color profiles similar to those in Kar 50 are not uncommon among Magellanic Irregular galaxies in the Virgo (e.g. Figures 8 and 9 of Binggeli & Cameron 1993) and M81 (Bremnes, Binggeli, & Prugniel 1998; Froebrich & Meusinger 2000) groups.

Bar instabilities can mix gas throughout a system (e.g. Friedli, Benz, & Kennicutt 1994), and produce flat light profiles. The disruption of bars may play an important role in the morphological evolution of dwarf galaxies, with the final system having a lower density

and different kinematical properties (Mayer et al. 2001). However, despite the flat light and color profiles of Kar 50, there is no evidence of a bar in Figure 1.

Most surveys of the structural characteristics of nearby dwarf systems have relied on observations at blue and visible wavelengths. The z' filter samples a wavelength interval that is less affected by recent star formation than filters covering shorter wavelengths, although the relatively blue $i' - z'$ color of Kar 50 suggests that a significant fraction of the light near $1\mu\text{m}$ still comes from a young component. JHK observations of Kar 50 will be of interest, as they will allow firmer constraints to be placed on (1) the size of any old stellar substrate, (2) the degree of central concentration of this component, and (3) the presence of a bar, given that bars are more easily detected in H than at optical wavelengths (Eskridge et al. 2000).

In the absence of subsequent star-forming episodes, Kar 50 will evolve into a very low surface brightness system. To estimate the surface brightness of Kar 50 after fading we adopt, based on the integrated colors of the galaxy, a $M/L = 0.5$ solar, and further assume that the final $M/L = 1.0$, based on the integrated color of a ‘typical’ low surface brightness system in the Virgo cluster, for which $\overline{B - V} = 0.6$ (e.g. Impey, Bothun, & Malin 1988). Kar 50 will then fade by 0.8 mag per square arcsec if the population ages passively, and the central surface brightness will drop to $\sim 25 \text{ mag arcsec}^{-2}$ in g' . This is within the range of central surface brightnesses seen among low surface brightness galaxies in Virgo, some of which also have flat central light profiles (e.g. Impey et al. 1988).

Sincere thanks are extended to Jean-Rene Roy and Sidney van den Bergh for discussions regarding bars and the morphological properties of dwarf irregular galaxies.

REFERENCES

- Barazza, F. D., Binggeli, B., & Prugniel, P. 2001, *A&A*, 373, 12 (BBP)
- Binggeli, B., & Cameron, L. M. 1993, *A&AS*, 98, 297
- Bremnes, T., Binggeli, B., & Prugniel, P. 1998, *A&A*, 129, 313
- Caldwell, N., Armandroff, T. E., Da Costa, G. S., & Seitzer, P. 1998, *AJ*, 115, 535
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245
- Crampton, D. et al. 2000, *SPIE*, 4008, 114
- Davidge, T. J. 1992, *AJ*, 103, 1259
- Davidge, T. J., Da Costa, G. S., Jorgensen, I., & Allington-Smith, J. R. 2002, *AJ*, in press.
- Dohm-Palmer, R. C. et al. 1998, *AJ*, 116, 1227
- Eskridge, P. B. et al. 2000, *AJ*, 119, 536
- Fan, X. 1999, *AJ*, 117, 2528
- Friedli, D., Benz, W., & Kennicutt, R. 1994, *ApJ*, 430, L105
- Froebrich, D., & Meusinger, H. 2000, *A&AS*, 145, 229
- Fukugita, M., Ichikawa, T., Gunn, J. E., Doi, M., Shimasaku, K., & Schneider, D. P. 1996, *AJ*, 111, 1748
- Humphreys, R. M., & McElroy, D. B. 1984, *ApJ*, 284, 565
- Impey, C., Bothun, G., & Malin, D. 1988, *ApJ*, 330, 634
- Karachentseva, V. E., Karachentsev, I. D., & Börngen, F. 1985, *A&AS*, 60, 213 (KKR)
- Karachentseva, V. E., Karachentsev, I. D., Richter, G. M., von Berlepsch, R., & Fritze, K. 1987, *AN*, 308, 247 (K87)
- Krisciunas, K., Margon, B., & Szkody, P. 1998, *PASP*, 110, 1342
- Lenz, D. D., Newberg, H. J., Rosner, R., Richards, G. T., & Stoughton, C. 1998, *ApJS*, 119, 121
- Mayer, L., Governato, F., Colpi, M., Moore, B., Quinn, T., Wadsley, J., Stadel, J., & Lake, G. 2001, *ApJ*, 547, L123
- O'Donnell, J. E. 1994, *ApJ*, 422, 1580
- Piotto, G., King, I. R., & Djorgovski, S. 1988, *AJ*, 96, 1918
- Rozanski, R., Rowan-Robinson, M. 1994, *MNRAS*, 271, 530
- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525

- Schneider, D. P., et al. 2001, *AJ*, 121, 1232
- Stetson, P. B. 1987, *PASP*, 99, 191
- Stetson, P. B., & Harris, W. E. 1988, *AJ*, 96, 909
- van den Bergh, S. 2000, *The Galaxies of the Local Group* (Cambridge: Cambridge University Press).
- Youngblood, A. J., & Hunter, D. A. 1999, *ApJ*, 519, 55

FIGURE CAPTIONS

Fig. 1.— The portion of the final r' image containing Kar 50. North is to the top and East is to the left. The square marks the HII region discussed in §4, while the two circles mark the intervals used in the stellar content investigation, which have outer radii of 22 and 31 arcsec.

Fig. 2.— The g' , r' , i' , and z' surface brightness profiles of Kar 50, obtained using the azimuthal filtering technique described in the text to suppress bright resolved stars. Note the remarkably flat profiles within 8 arcsec of the galaxy center. The error bars show the uncertainty in the largest radii measurements due to uncertainties in the sky level.

Fig. 3.— The $(g' - i')$ and $(r' - z')$ color profiles of Kar 50, corrected for line of sight reddening. The error bars show the uncertainty in the largest radii measurements due to uncertainties in the sky level, and the errors in the colors become progressively smaller than this with decreasing radius. The dashed lines show colors predicted for various effective temperatures from the $\log(g) = 4.5$ solar metallicity models of Lenz et al. (1998). Note that there is no evidence for significant gradients within 15 arcsec of the galaxy center. While apparent departures from a flat color profile occur at larger radii, these are not significant given the uncertainty in the colors at these radii.

Fig. 4.— The $(r', g' - r')$ CMDs of objects within 22 arcsec of the center of Kar 50 (left hand panel), and those located between 22 and 31 arcsec from the galaxy center (right hand panel).

Fig. 5.— The $(i', r' - i')$ CMDs of objects within 22 arcsec of the center of Kar 50 (left hand panel), and those located between 22 and 31 arcsec from the galaxy center (right hand panel).

Fig. 6.— The $(g' - r', r' - i')$ TCDs of objects within 22 arcsec of the center of Kar 50 (left hand panel), and those located between 22 and 31 arcsec from the galaxy center (right hand panel). The solid line shows the locus of the solar metallicity $\log(g)=4.5$ models of Lenz et al. (1998), while the dashed line is the SDSS stellar sequence from Figure 3 of Schneider et al. (2001). The reddening vector follows the relations listed in Table 6 of Schlegel et al. (1998), which in turn is based on the Cardelli, Clayton, & Mathis (1989) and O'Donnell (1994) reddening curves. The length of the reddening vector corresponds to $A_{g'} = 1$ mag. Note that the majority of sources within 22 arcsec of the center of Kar 50 have $g' - r' < 0$, whereas those in the background field have $g' - r' > 0$.

Fig. 7.— The $(r', g' - r')$ and $(i', r' - i')$ CMDs of stars within 22 arcsec of the center of

Kar 50. The data have been corrected for line of sight extinction predicted from the Schlegel et al. (1998) maps. The dashed line marks $g' - r' = 0$, which is the criterion adopted for rejecting background objects; sources falling to the right of this line in the $(r', g' - r')$ CMD have a high likelihood of being contaminating objects. The solid squares in the right hand panel mark those sources detected in g', r' , and i' having $g' - r' > 0$, and as such are probable foreground stars or background galaxies. The point corresponding to the candidate HII region is also indicated in each CMD.

Fig. 8.— The $(r', g' - r')$ CMD of stars within 22 arcsec of the center of Kar 50. The data have been corrected for line of sight extinction as predicted from the Schlegel et al. (1998) maps. The dashed line marks $g' - r' = 0$; sources falling to the right of this line are likely unrelated to Kar 50. The source identified as an HII region is labelled, as are the loci of supergiants and upper main sequence stars in the Milky-Way and the Magellanic Clouds, assuming a distance modulus of 30.4.

Quantity	Value	Reference
RA (1950)	$07^h48^m25^s$	KKR
Dec (1950)	$61^\circ31'08''$	KKR
l_{II} (1950)	155.4789	
b_{II} (1950)	30.8665	
Type	dImV	KKR
B_T	17.88	BBP
r_{eff}^B	11.1 arcsec	BBP
B–V	0.24	BBP
E(B–V)	0.05	Schlegel et al. (1998)

Table 1: Basic properties of Kar 50

Filter	Date (UT)	Exposure time (sec)	FWHM (arcsec)
g'	Nov 22, 2001	2×900	1.00
r'	Nov 19, 2001	6×400	0.65
i'	Nov 19, 2001	6×400	0.65
z'	Nov 22, 2001	12×200	0.85

Table 2: Details of Observations

r'_0	$(g' - r')_0$	$(B - V)_0^a$	V_0^a	B_0
22.76	−0.19	−0.01	22.66	22.65
23.57	−0.18	0.00	23.47	23.46
23.58	−0.21	−0.03	23.46	23.43

Table 3: Photometric Properties of the Three Brightest Stars

^aComputed from r'_0 and $(g' - r')_0$ using the relations given in Section 2.1 of Krisciunas et al. (1998).















